

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 7/50	A1	(11) International Publication Number: WO 97/03648 (43) International Publication Date: 6 February 1997 (06.02.97)
(21) International Application Number: PCT/US96/11789 (22) International Filing Date: 17 July 1996 (17.07.96) (30) Priority Data: 08/506,149 24 July 1995 (24.07.95) US (71) Applicant: THE PROCTER & GAMBLE COMPANY [US/US]; One Procter & Gamble Plaza, Cincinnati, OH 45202 (US). (72) Inventors: McATEE, David, Michael; 100 Bent Tree Drive, Fairfield, OH 45019 (US). ALBACARYS, Lourdes, Dessus; 5819 Owl Nest Drive, West Chester, OH 45069 (US). LISTRO, Joseph, Anthony; 9758 Tulip Tree Court, Love- land, OH 45140 (US). (74) Agents: REED, T., David et al.; The Procter & Gamble Company, 5299 Spring Grove Avenue, Cincinnati, OH 45217 (US).	(81) Designated States: AU, CA, CN, CZ, JP, KR, MX, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the</i> <i>claims and to be republished in the event of the receipt of</i> <i>amendments.</i>	
(54) Title: COMPOSITIONS FOR TOPICAL DELIVERY OF ACTIVE INGREDIENTS (57) Abstract <p>The compositions of the present invention are useful for the topical delivery of a wide variety of active ingredients. These compositions are particularly useful for treating conditions such as acne and its attendant skin lesions, blemishes, and other imperfections. These compositions are nonirritating to the skin and also provide skin feel benefits. These compositions can be in the form of leave-on products and products that are rinsed or wiped from the skin after use.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MZ	Mozambique	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

COMPOSITIONS FOR TOPICAL DELIVERY OF ACTIVE INGREDIENTS

TECHNICAL FIELD

5 The compositions of the present invention are useful for the topical delivery of a wide variety of active ingredients. These compositions are particularly useful for treating conditions such as acne and its attendant skin lesions, blemishes, and other imperfections. These compositions are useful for clarifying the skin and reducing pore size. These compositions are nonirritating to the skin and also provide improved skin feel benefits. These compositions can be in the form of leave-
10 on products and products that are rinsed or wiped from the skin after use.

BACKGROUND OF THE INVENTION

 The treatment of human skin with various agents has been undertaken for many years with the goal being to keep the skin in a smooth and supple condition. Skin has the tendency to dry out when exposed to low humidity or to harsh detergent solutions for extended periods of time. From a
15 physiological standpoint, dryness is a measure of the water content of the skin. Under normal conditions, the water content and vapor pressure of the epidermis are higher than those of the surrounding air, with consequent evaporation of water from the skin surface. Skin becomes dry because of excessive loss of water from its surface which results in loss of water from the stratum corneum. Low humidity speeds up this process, exacerbating the drying of skin.

20 Continuous and prolonged immersion in soap or detergent solutions can contribute to dryness of the stratum corneum. The reason for this is that the surfactant medium promotes dissolution of the skin surface and horny layer lipids, and the dissolution of the hygroscopic water-soluble components in the skin.

 Also, in normal skin, the stratum corneum is shed as individual cells or as small clusters of
25 cells. Skin problems such as dry skin, psoriasis, ichthyosis, dandruff, callus, photodamaged skin, aged skin, and sunburn can be described as disorders of keratinization in which the shedding of stratum corneum cells at the skin surface is altered relative to normal, young, healthy skin. Such alteration results in shedding of large clusters of cells leading to visible scaling of the skin, a build-up of keratinaceous material on the surface or in follicles or ducts, and a rough texture to the skin
30 surface. These conditions can be improved by removal of the outermost keratinaceous material. In other words, by desquamation.

 Acne is a condition of the human skin characterized by an excess flow of sebum, or skin oil, from the sebaceous glands located in the pilosebaceous apparatus. Sebum reaches the skin surface through the duct of the hair follicle. The presence of excessive amounts of sebum in the duct and on
35 the skin acts to block or stagnate the continuous flow of sebum from the follicular duct, thus producing a thickening and a solidification of the sebum to form a solid plug known as a comedone.

When this process occurs, hyperkeratinization of the follicular opening is stimulated, thus completely closing the duct. The usual results are papules, pustules, or cysts, often contaminated with bacteria which cause secondary infections. Acne is particularly characterized by the presence of comedones, inflammatory papules, pustules, or cysts. The effect of acne ranges from slight skin
5 irritation and pitting to disfiguring scars.

Many topical therapeutic agents are employed in the treatment of acne to prevent the blocking of the follicular duct, to reopen the duct once it has become blocked, to act against the infecting bacteria or the thickened sebum, and to provide combinations of each of these actions. The horny outer layer of the skin, which is known as the stratum corneum, is formed of dead cells
10 composed largely of keratin. Therapeutic agents which act to prevent the blocking of the follicular duct by promoting the removal or sloughing off of excess keratin are known as keratolytic agents. However, many anti-acne compositions are harsh and irritating to the skin.

Therefore, there is an ongoing need to effectively deliver a wide variety of active ingredients, particularly anti-acne actives, to the skin, either via direct application of such a composition, or in
15 the case of a cleansing composition, via the cleansing process. There is a need for developing products which are gentle and nonirritating to the skin.

It has been found in the present invention that skin care compositions containing a combination of an amphoteric surfactant and an anionic surfactant, further in combination with an active ingredient, are useful for providing these skin care benefits. When the active ingredient is
20 salicylic acid or benzoyl peroxide, these compositions have been found especially useful for treating or cleansing the skin, especially skin that is afflicted with acne. These compositions are found to provide improved anti-acne efficacy. These compositions have been found to be gentle and nonirritating and to leave the skin feeling soft and smooth.

It is therefore an object of the present invention to provide skin care compositions for topical
25 application to the skin.

It is another object of the present invention to provide skin care compositions which are useful for efficiently delivering a wide variety of active ingredients to the skin.

It is another object of the present invention to provide skin care compositions having improved skin conditioning and dryness reducing properties.

30 It is another object of the present invention to provide compositions which are useful for treating and preventing acne, while being mild and nonirritating to the skin.

It is another object of the present invention to provide skin care compositions which, when in the form of cleansing compositions, are useful for delivering a wide variety of active ingredients to the skin via the cleansing process.

35 It is another object of the present invention to provide methods for treating the skin.

It is another object of the present invention to provide methods for delivering active ingredients to the skin, to provide methods for treating acne, to provide methods for treating dry


skin, to provide methods for cleansing the skin, to provide methods for reducing pore size, to provide methods for clarifying the skin, and to provide methods for desquamating the skin.

These and other objects will become readily apparent from the detailed description which follows.

SUMMARY OF THE INVENTION

The present invention relates to topical personal care compositions comprising:

(a) from about 0.1% to about 20% by weight of an amphoteric surfactant having the following structure




wherein R^1 is unsubstituted, saturated or unsaturated, straight or branched chain alkyl having from about 9 to about 22 carbon atoms; m is an integer from 1 to about 3; n is 0 or 1; R^2 and R^3 are independently selected from alkyl having from 1 to about 3 carbon atoms and monohydroxyalkyl having from 1 to about 3 carbon atoms; R^4 is selected from saturated or unsaturated alkyl having from 1 to about 5 carbon atoms and saturated or unsaturated monohydroxyalkyl having from 1 to about 5 carbon atoms; X is selected from the group consisting of CO_2 , SO_3 , and SO_4 ; and pharmaceutically acceptable salts of the foregoing compounds;

- (b) from about 0.1% to about 20% by weight of an anionic surfactant,
- (c) from about 0.001% to about 20% of an active ingredient, and
- (d) from about 40% to about 99.799% by weight water.

The present invention also relates to methods for treating the skin, specifically to methods of treating conditions such as acne, and to methods for conditioning, cleansing, clarifying, reducing pore size, and desquamating the skin, utilizing these foregoing compositions.

The present invention also relates to methods of preparing a composition comprising

(a) from about 0.1% to about 20% by weight of an amphoteric surfactant having the following structure



wherein R^1 is unsubstituted, saturated or unsaturated, straight or branched chain alkyl having from about 9 to about 22 carbon atoms; m is an integer from 1 to about 3; n is 0 or 1; R^2 and R^3 are

- independently selected from alkyl having from 1 to about 3 carbon atoms and monohydroxyalkyl having from 1 to about 3 carbon atoms; R^4 is selected from saturated or unsaturated alkyl having from 1 to about 5 carbon atoms and saturated or unsaturated monohydroxyalkyl having from 1 to about 5 carbon atoms; X is selected from the group consisting of CO_2 , SO_3 , and SO_4 ; and
- 5 pharmaceutically acceptable salts of the foregoing compounds;
- (b) from about 0.1% to about 20% by weight of an anionic surfactant,
 - (c) from about 0.001% to about 20% of an active ingredient, and
 - (d) from about 40% to about 99.799% by weight water,

comprising the steps of:

- 10 (i) combining an aqueous solution of said amphoteric surfactant and an aqueous solution of said anionic surfactant to form an aqueous dispersion of a complex of said amphoteric and said anionic surfactant, and
- (ii) combining said aqueous dispersion of said complex with said active ingredient.

15 All percentages and ratios used herein are by weight of the total composition and all measurements made are at 25°C or room temperature, unless otherwise designated. All weight percentages, unless otherwise indicated, are on an active weight basis. The invention hereof can comprise, consist of, or consist essentially of, the essential as well as the optional ingredients and additional components described herein.

DETAILED DESCRIPTION OF THE INVENTION

20 The compositions of the present invention are useful for delivering a wide variety of active ingredients to the surface of the skin. These compositions, when they contain anti-acne actives provide improved anti-acne efficacy versus conventional formulas.

Without being limited by theory it is believed that the amphoteric surfactant of these compositions complexes with the anionic surfactant. This postulated complex is believed to be

25 highly stable relative to the individual surfactant components. This complex is useful for aiding in the delivery of the active ingredients to the skin, thereby aiding their efficacy. In the case of a cleansing composition, this complex tends to deposit out from the composition thereby helping to carry the active ingredients to the skin's surface during the cleansing process. Because the postulated complex contains an amphoteric and an anionic surfactant, this complex is also effective

30 for cleansing the skin and for promoting the desquamation process. Because the charges on the individual surfactants are complexed, the surfactants are rendered less harsh and irritating to the skin versus the free surfactants.

The compositions of the present invention can be formulated into a wide variety of product types including, but not limited to creams, lotions, mousses, sprays, "rinse-off" cleansers, "water-

35 less" cleansers, bars, gels, and the like. The term "rinse-off," as used herein, means that the composition is in a form that can be used in a cleansing process whereby the composition is ultimately rinsed or washed from the skin with water to complete the cleansing process. The term

"water-less," as used herein, means that the composition is in a form that can be used in a cleansing process without water whereby the composition is typically removed by wiping with a device such as a cotton ball, a cotton pad, a tissue, a towel, and the like.

The term "pharmaceutically-acceptable," as used herein, means that the compositions and components thereof so described are of sufficiently high purity and are suitable for use in contact with human skin and tissues without undue toxicity, irritation, incompatibility, instability, allergic response, and the like.

The term "pharmaceutically-acceptable salts," as used herein means any of the commonly-used salts that are suitable for use in contact with human skin and tissues without undue toxicity, irritation, incompatibility, instability, allergic response, and the like.

AMPHOTERIC SURFACTANT

The compositions of the present invention comprise from about 0.1% to about 20%, more preferably from about 0.2% to about 10%, and most preferably from about 0.5% to about 5% of an amphoteric surfactant.

The term "amphoteric surfactant," as used herein, is also intended to encompass zwitterionic surfactants, which are well known to formulators skilled in the art as a subset of amphoteric surfactants.

A wide variety of amphoteric surfactants can be used in the compositions of the present invention. Particularly useful are those which are broadly described as derivatives of aliphatic secondary and tertiary amines, preferably wherein the nitrogen is in a cationic state, in which the aliphatic radicals can be straight or branched chain and wherein one of the radicals contains an ionizable water solubilizing group, e.g., carboxy, sulfonate, sulfate, phosphate, or phosphonate.


Nonlimiting examples of amphoteric surfactants useful in the compositions of the present invention are disclosed in McCutcheon's, Detergents and Emulsifiers, North American edition (1986), published by Allured Publishing Corporation; and McCutcheon's, Functional Materials, North American Edition (1992); both of which are incorporated by reference herein in their entirety.

Preferred amphoteric or zwitterionic surfactants are the betaines, sultaines, and hydroxysultaines. Examples of betaines include the higher alkyl betaines, such as coco dimethyl carboxymethyl betaine, lauryl dimethyl carboxymethyl betaine, lauryl dimethyl alphacarboxyethyl betaine, cetyl dimethyl carboxymethyl betaine, cetyl dimethyl betaine (available as Lonzaine 16SP from Lonza Corp.), lauryl bis-(2-hydroxyethyl) carboxymethyl betaine, stearyl bis-(2-hydroxypropyl) carboxymethyl betaine, oleyl dimethyl gamma-carboxypropyl betaine, lauryl bis-(2-hydroxypropyl)alpha-carboxyethyl betaine, coco dimethyl sulfopropyl betaine, stearyl dimethyl sulfopropyl betaine, stearyl betaine, lauryl dimethyl sulfoethyl betaine, lauryl bis-(2-hydroxyethyl) sulfopropyl betaine, and amidobetaines and amidosulfobetaines (wherein the $RCONH(CH_2)_3$ radical is attached to the nitrogen atom of the betaine), oleyl betaine (available as

amphoteric Velvetex OLB-50 from Henkel), and cocamidopropyl betaine (available as Velvetex BK-35 and BA-35 from Henkel).

Examples of sultaines and hydroxysultaines include materials such as cocamidopropyl hydroxysultaine (available as Mirataine CBS from Rhone-Poulenc).


5 Preferred for use herein are amphoteric surfactants having the following structure:




10 wherein R^1 is unsubstituted, saturated or unsaturated, straight or branched chain alkyl having from about 9 to about 22 carbon atoms. Preferred R^1 has from about 11 to about 18 carbon atoms; more preferably from about 12 to about 18 carbon atoms; more preferably still from about 14 to about 18 carbon atoms; m is an integer from 1 to about 3, more preferably from about 2 to about 3, and more preferably about 3; n is either 0 or 1, preferably 1; R^2 and R^3 are independently selected from the group consisting of alkyl having from 1 to about 3 carbon atoms, unsubstituted or mono-substituted with hydroxy, preferred R^2 and R^3 are CH_3 ; X is selected from the group consisting of CO_2 , SO_3 and SO_4 ; R^4 is selected from the group consisting of saturated or unsaturated, straight or branched chain alkyl, unsubstituted or monosubstituted with hydroxy, having from 1 to about 5 carbon atoms. When X is CO_2 , R^4 preferably has 1 or 3 carbon atoms, more preferably 1 carbon atom. When X is SO_3 or SO_4 , R^4 preferably has from about 2 to about 4 carbon atoms, more preferably 3 carbon atoms.

20 Examples of amphoteric surfactants of the present invention include the following compounds:

Cetyl dimethyl betaine (this material also has the CTFA designation cetyl betaine)




Cocamidopropylbetaine




wherein R has from about 9 to about 13 carbon atoms

Cocamidopropyl hydroxy sultaine




5 wherein R has from about 9 to about 13 carbon atoms,

Stearyl dimethyl betaine



10 and, Behenyl Betaine



Preferred amphoteric surfactants of the present invention include cetyl dimethyl betaine, cocoamidopropyl betaine, stearyl dimethyl betaine, and cocoamidopropyl hydroxy sultaine. Still
15 more preferred are cetyl dimethyl betaine, stearyl dimethyl betaine, and cocamidopropyl betaine. Most preferred is cetyl dimethyl betaine.

Examples of other useful amphoteric surfactants are alkyliminoacetates, and iminodialkanoates and aminoalkanoates of the formulas $\text{RN}(\text{CH}_2)_m\text{CO}_2\text{M}]_2$ and $\text{RNH}(\text{CH}_2)_m\text{CO}_2\text{M}$ wherein m is from 1 to 4, R is a C₈-C₂₂ alkyl or alkenyl, and M is H, alkali
20 metal, alkaline earth metal ammonium, or alkanolammonium. Also included are imidazolinium and ammonium derivatives. Specific examples of suitable amphoteric surfactants include sodium 3-dodecyl-aminopropionate, sodium 3-dodecylaminopropane sulfonate, N-alkyltaurines such as the one prepared by reacting dodecylamine with sodium isethionate according to the teaching of U.S. Patent 2,658,072 which is incorporated herein by reference in its entirety; N-higher alkyl aspartic
25 acids such as those produced according to the teaching of U.S. Patent 2,438,091 which is incorporated herein by reference in its entirety; and the products sold under the trade name "Miranol" and described in U.S. Patent 2,528,378, which is incorporated herein by reference in its entirety. Other examples of useful amphoterics include phosphates, such as coamidopropyl PG-dimonium chloride phosphate (commercially available as Monaquat PTC, from Monâ Corp.).


30 ANIONIC SURFACTANT

The compositions of the present invention comprise from about 0.1% to about 20%, more preferably from about 0.2% to about 10%, and most preferably from about 0.5% to about 5% of an anionic surfactant.

Nonlimiting examples of anionic surfactants useful in the compositions of the present invention are disclosed in McCutcheon's, Detergents and Emulsifiers, North American edition (1986), published by allured Publishing Corporation; McCutcheon's, Functional Materials, North American Edition (1992); and U.S. Patent No. 3,929,678, to Laughlin et al., issued December 30, 1975 all of which are incorporated by reference herein in their entirety.

A wide variety of anionic surfactants are useful herein. Nonlimiting examples of anionic surfactants include the alkoyl isethionates, and the alkyl and alkyl ether sulfates. The alkoyl isethionates typically have the formula $\text{RCO-OCH}_2\text{CH}_2\text{SO}_3\text{M}$ wherein R is alkyl or alkenyl of from about 10 to about 30 carbon atoms, and M is a water-soluble cation such as ammonium, sodium, potassium and triethanolamine. Nonlimiting examples of these isethionates include those alkoyl isethionates selected from the group consisting of ammonium cocoyl isethionate, sodium cocoyl isethionate, sodium lauroyl isethionate, sodium stearoyl isethionate, and mixtures thereof.

The alkyl and alkyl ether sulfates typically have the respective formulae ROSO_3M and $\text{RO}(\text{C}_2\text{H}_4\text{O})_x\text{SO}_3\text{M}$, wherein R is alkyl or alkenyl of from about 10 to about 30 carbon atoms, x is from about 1 to about 10, and M is a water-soluble cation such as ammonium, sodium, potassium and triethanolamine. Another suitable class of anionic surfactants are the water-soluble salts of the organic, sulfuric acid reaction products of the general formula:



wherein R_1 is chosen from the group consisting of a straight or branched chain, saturated aliphatic hydrocarbon radical having from about 8 to about 24, preferably about 10 to about 16, carbon atoms; and M is a cation. Still other anionic synthetic surfactants include the class designated as succinamates, olefin sulfonates having about 12 to about 24 carbon atoms, and b-alkyloxy alkane sulfonates. Examples of these materials are sodium lauryl sulfate and ammonium lauryl sulfate.

Other anionic materials include the sarcosinates nonlimiting examples of which include sodium lauroyl sarcosinate, sodium cocoyl sarcosinate, and ammonium lauroyl sarcosinate.

Other anionic materials useful herein are soaps (i.e. alkali metal salts, e.g., sodium or potassium salts) of fatty acids, typically having from about 8 to about 24 carbon atoms, preferably from about 10 to about 20 carbon atoms. The fatty acids used in making the soaps can be obtained from natural sources such as, for instance, plant or animal-derived glycerides (e.g., palm oil, coconut oil, soybean oil, castor oil, tallow, lard, etc.) The fatty acids can also be synthetically prepared. Soaps are described in more detail in U.S. Patent No. 4,557,853, cited above.

Other anionic materials include phosphates such as monoalkyl, dialkyl, and trialkylphosphate salts.

Other anionic materials include alkanoyl sarcosinates corresponding to the formula $RCON(CH_3)CH_2CH_2CO_2M$ wherein R is alkyl or alkenyl of about 10 to about 20 carbon atoms, and M is a water-soluble cation such as ammonium, sodium, potassium and trialkanolamine (e.g., triethanolamine), a preferred example of which is sodium lauroyl sarcosinate.

5 Nonlimiting examples of preferred anionic surfactants useful herein include those selected from the group consisting of sodium lauryl sulfate, ammonium lauryl sulfate, ammonium cetyl sulfate, sodium cetyl sulfate, sodium stearyl sulfate, ammonium cocoyl isethionate, sodium lauroyl isethionate, sodium lauroyl sarcosinate, and mixtures thereof.

Especially preferred for use herein is sodium lauryl sulfate.

10 ACTIVE INGREDIENTS

The compositions of the present invention comprise a safe and effective amount of one or more active ingredients or pharmaceutically-acceptable salts thereof.

The term "safe and effective amount" as used herein, means an amount of an active ingredient high enough to modify the condition to be treated or to deliver the desired skin benefit,
15 but low enough to avoid serious side effects, at a reasonable benefit to risk ratio within the scope of sound medical judgement. What is a safe and effective amount of the active ingredient will vary with the specific active, the ability of the active to penetrate through the skin, the age, health condition, and skin condition of the user, and other like factors.

Typically, the active ingredients of the present invention comprise from about 0.001% to
20 about 20%, preferably from about 0.01% to about 15%, and more preferably from about 0.025% to about 10% by weight of the composition.

The active ingredients useful herein can be categorized by their therapeutic benefit or their postulated mode of action. However, it is to be understood that the active ingredients useful herein can in some instances provide more than one therapeutic benefit or operate via more than one mode
25 of action. Therefore, classifications herein are made for the sake of convenience and are not intended to limit the active ingredient to that particular application or applications listed. Also, pharmaceutically-acceptable salts of these active ingredients are useful herein. The following active ingredients are useful in the compositions of the present invention.

Anti-Acne Actives: Examples of useful anti-acne actives include the keratolytics such as salicylic acid (o-hydroxybenzoic acid), derivatives of salicylic acid such as 5-octanoyl salicylic acid, and
30 resorcinol; retinoids such as retinoic acid and its derivatives (e.g., cis and trans); sulfur-containing D and L amino acids and their derivatives and salts, particularly their N-acetyl derivatives, a preferred example of which is N-acetyl-L-cysteine; lipoic acid; antibiotics and antimicrobials such as benzoyl peroxide, octopirox, tetracycline, 2,4,4'-trichloro-2'-hydroxy diphenyl ether, 3,4,4'-
35 trichlorobanilide, azelaic acid and its derivatives, phenoxyethanol, phenoxypropanol, phenoxyisopropanol, ethyl acetate, clindamycin and meclocycline; sebstats such as flavonoids; and bile salts such as scymnol sulfate and its derivatives, deoxycholate, and cholate.

Anti-Wrinkle and Anti-Skin Atrophy Actives: Examples of antiwrinkle and anti-skin atrophy actives include retinoic acid and its derivatives (e.g., cis and trans); retinol; retinyl esters; salicylic acid and derivatives thereof; sulfur-containing D and L amino acids and their derivatives and salts, particularly the N-acetyl derivatives, a preferred example of which is N-acetyl-L-cysteine; thiols, eg. ethane thiol; alpha-hydroxy acids, e.g. glycolic acid, and lactic acid; phytic acid, lipoic acid; lysophosphatidic acid, and skin peel agents (e.g., phenol and the like).

Non-Steroidal Anti-Inflammatory Actives (NSAIDS): Examples of NSAIDS include the following categories: propionic acid derivatives; acetic acid derivatives; fenamic acid derivatives; biphenylcarboxylic acid derivatives; and oxicams. All of these NSAIDS are fully described in U.S. Patent 4,985,459 to Sunshine et al., issued January 15, 1991, incorporated by reference herein in its entirety. Examples of useful NSAIDS include acetyl salicylic acid, ibuprofen, naproxen, benoxaprofen, flurbiprofen, fenoprofen, fenbufen, ketoprofen, indoprofen, piroprofen, carprofen, oxaprozin, pranoprofen, miroprofen, tioxaprofen, suprofen, alminoprofen, tiaprofenic acid, fluprofen and bucloxic acid. Also useful are the steroidal anti-inflammatory drugs including hydrocortisone and the like.

Topical Anesthetics: Examples of topical anesthetic drugs include benzocaine, lidocaine, bupivacaine, chlorprocaine, dibucaine, etidocaine, mepivacaine, tetracaine, dyclonine, hexylcaine, procaine, cocaine, ketamine, pramoxine, phenol, and pharmaceutically acceptable salts thereof.

Antimicrobial and Antifungal Actives: Examples of antimicrobial and antifungal actives include β -lactam drugs, quinolone drugs, ciprofloxacin, norfloxacin, tetracycline, erythromycin, amikacin, 2,4,4'-trichloro-2'-hydroxy diphenyl ether, 3,4,4'-trichlorobanilide, phenoxyethanol, phenoxy propanol, phenoxyisopropanol, doxycycline, capreomycin, chlorhexidine, chlortetracycline, oxytetracycline, clindamycin, ethambutol, hexamidine isethionate, metronidazole, pentamidine, gentamicin, kanamycin, lineomycin, methacycline, methenamine, minocycline, neomycin, netilmicin, paromomycin, streptomycin, tobramycin, miconazole, tetracycline hydrochloride, erythromycin, zinc erythromycin, erythromycin estolate, erythromycin stearate, amikacin sulfate, doxycycline hydrochloride, capreomycin sulfate, chlorhexidine gluconate, chlorhexidine hydrochloride, chlortetracycline hydrochloride, oxytetracycline hydrochloride, clindamycin hydrochloride, ethambutol hydrochloride, metronidazole hydrochloride, pentamidine hydrochloride, gentamicin sulfate, kanamycin sulfate, lineomycin hydrochloride, methacycline hydrochloride, methenamine hippurate, methenamine mandelate, minocycline hydrochloride, neomycin sulfate, netilmicin sulfate, paromomycin sulfate, streptomycin sulfate, tobramycin sulfate, miconazole hydrochloride, amantadine hydrochloride, amantadine sulfate, octopirox, parachlorometa xyleneol, nystatin, tolnaftate and clotrimazole.

Preferred examples of actives useful herein include those selected from the group consisting of salicylic acid, benzoyl peroxide, 3-hydroxy benzoic acid, glycolic acid, lactic acid, 4-hydroxy benzoic acid, acetyl salicylic acid, 2-hydroxybutanoic acid, 2-hydroxypentanoic acid, 2-

hydroxyhexanoic acid, cis-retinoic acid, trans-retinoic acid, retinol, phytic acid, N-acetyl-L-cysteine, lipoic acid, azelaic acid, arachidonic acid, benzoylperoxide, tetracycline, ibuprofen, naproxen, hydrocortisone, acetaminophen, resorcinol, phenoxyethanol, phenoxypropanol, phenoxyisopropanol, 2,4,4'-trichloro-2'-hydroxy diphenyl ether, 3,4,4'-trichlorocarbaniide, octopirox, lidocaine
5 hydrochloride, clotrimazole, miconazole, neocycin sulfate, and mixtures thereof.

More preferred examples of actives useful herein include those selected from the group consisting of salicylic acid, acetyl salicylic acid, cis-retinoic acid, trans-retinoic acid, retinol, phytic acid, N-acetyl-L-cysteine, lipoic acid, azelaic acid, tetracycline, ibuprofen, naproxen, acetaminophen, hydrocortisone, resorcinol, phenoxyethanol, phenoxypropanol, phenoxyisopropanol,
10 2,4,4'-trichloro-2'-hydroxy diphenyl ether, 3,4,4'-trichlorocarbaniide, octopirox, and mixtures thereof.

More preferred examples of actives useful herein include those selected from the group consisting of salicylic acid, benzoyl peroxide, cis-retinoic acid, trans-retinoic acid, retinol, phytic acid, N-acetyl L-cysteine, azelaic acid, lipoic acid, resorcinol, lactic acid, glycolic acid, ibuprofen, naproxen, hydrocortisone, phenoxyethanol, phenoxypropanol, phenoxyisopropanol, 2,4,4'-trichloro-
15 2'-hydroxy diphenyl ether, 3,4,4'-trichlorocarbaniide, and mixtures thereof.

Most preferred are salicylic acid and benzoyle peroxide.

WATER

The compositions of the present invention comprise from about 40% to about 99.799%, more preferably from about 60% to about 95%, and most preferably from about 70% to about 90% of water. The exact level of water will depend upon the form of the product and the desired moisture content.
20

ADDITIONAL COMPONENTS

The compositions of the present invention can comprise a wide range of additional components. The CTFA Cosmetic Ingredient Handbook, Second Edition, 1992, which is incorporated by reference herein in its entirety, describes a wide variety of cosmetic and pharmaceutical ingredients commonly used in the skin care industry, which are suitable for use in the compositions of the present invention. Nonlimiting examples of functional classes of ingredients are described at page 537 of this reference. Examples of these functional classes
25 include: absorbents, abrasives, anti-acne agents, anticaking agents, antifoaming agents, antimicrobial agents, antioxidants, binders, biological additives, buffering agents, bulking agents, chelating agents, chemical additives, colorants, cosmetic astringents, cosmetic biocides, denaturants, drug astringents, external analgesics, film formers, fragrance components, humectants, opacifying agents, pH adjusters, plasticizers, preservatives, propellants, reducing agents, skin
30 bleaching agents, skin-conditioning agents (emollient, humectants, miscellaneous, and occlusive), skin protectants, solvents, foam boosters, hydrotropes, solubilizing agents, suspending agents (nonsurfactant), sunscreen agents, ultraviolet light absorbers, and viscosity increasing agents
35

(aqueous and nonaqueous). Examples of other functional classes of materials useful herein that are well known to one of ordinary skill in the art include emulsifiers, sequestrants, skin sensates, and the like.

Nonlimiting examples of these additional components cited in the CTFA Cosmetic Ingredient Handbook as well as other materials useful herein, include the following: vitamins and derivatives thereof [e.g., vitamin C, Vitamin A (i.e. retinoic acid), retinol, retinoids, panthenol, panthehol esters, tocopherol, tocopherol esters, and the like]; oil or sebum control agents such as clays silicones and drug actives; suncreening agents; other silicone materials such as dimethiconol, dimethicone copolyol, and amodimethicone, and the like); anti-oxidants; anti-microbial agents; preservatives; emulsifiers; polyethyleneglycols and polypropyleneglycols; polymers for aiding the film-forming properties and substantivity of the composition (such as a copolymer of eicosene and vinyl pyrrolidone, an example of which is available from GAF Chemical Corporation as Ganex[®] V-220); preservatives for maintaining the antimicrobial integrity of the compositions; anti-acne medicaments (e.g., resorcinol, sulfur, salicylic acid, erythromycin, zinc, and the like); skin bleaching (or lightening) agents including but not limited to hydroquinone, kojic acid; antioxidants; chelators and sequestrants; thickening agents such as carbomers (homopolymers of acrylic acid crosslinked with an allyl ether of pentaerythritol or an allyl ether of sucrose), crosslinked and noncrosslinked nonionic and cationic polyacrylamides [e.g., Salcare[®] SC92 which has the CTFA designation polyquaternium 32 (and) mineral oil, and Salcare[®] SC 95 which has the CTFA designation polyquaternium 37 (and) mineral oil (and) PPG-1 trideceth-6, and the nonionic Seppi-Gel polyacrylamides available from Seppic Corp.]; proteins and peptides; enzymes; ceramides; aesthetic components such as fragrances, pigments, colorings, essential oils, skin senates, astringents, skin soothing agents, skin healing agents and the like, [nonlimiting examples of these aesthetic components include clove oil, menthol, camphor, eucalyptus oil, eugenol, menthyl lactate, witch hazel distillate, bisabolol, dipotassium glycyrrhizinate and the like]; and skin conditioning agents such as urea and glycerol, and also the propoxylated glycerols described in U.S. Patent No. 4,976,953, to Orr et al., issued December 11, 1990, which is incorporated by reference herein in its entirety.

Some of these additional ingredients are described in more detail below.

30 ADDITIONAL SURFACTANTS


The compositions of the present invention, in addition to the required surfactant materials, can comprise additional surfactant materials. Especially useful are cationic and nonionic surfactants.

The compositions of the present invention can comprise from about 0.1% to about 15%, more preferably from about 0.2% to about 10%, and most preferably from about 0.5% to about 5% of a cationic surfactant.

Nonlimiting examples of cationic surfactants useful in the compositions of the present invention are disclosed in McCutcheon's, Detergents and Emulsifiers, North American edition (1986), published by allured Publishing Corporation; and McCutcheon's, Functional Materials, North American Edition (1992); both of which are incorporated by reference herein in their entirety. Without being limited by theory, it is believed that such cationic materials can also provide an antimicrobial effect to the compositions herein. Therefore, cationic materials having antimicrobial properties are highly useful herein.

Nonlimiting examples of cationic surfactants useful herein include cationic ammonium salts such as those having the formula:

10



wherein R_1 is selected from an alkyl group having from about 12 to about 22 carbon atoms, or aromatic, aryl or alkaryl groups having from about 12 to about 22 carbon atoms; R_2 , R_3 , and R_4 are independently selected from hydrogen, an alkyl group having from about 1 to about 22 carbon atoms, or aromatic, aryl or alkaryl groups having from about 12 to about 22 carbon atoms; and X is an anion selected from chloride, bromide, iodide, acetate, phosphate, nitrate, sulfate, methyl sulfate, ethyl sulfate, tosylate, lactate, citrate, glycolate, and mixtures thereof. Additionally, the alkyl groups can also contain ether linkages, or hydroxy or amino group substituents (e.g., the alkyl groups can contain polyethylene glycol and polypropylene glycol moieties).

More preferably, R_1 is an alkyl group having from about 12 to about 22 carbon atoms; R_2 is selected from H or an alkyl group having from about 1 to about 22 carbon atoms; R_3 and R_4 are independently selected from H or an alkyl group having from about 1 to about 3 carbon atoms; and X is as described in the previous paragraph.

Most preferably, R_1 is an alkyl group having from about 12 to about 22 carbon atoms; R_2 , R_3 , and R_4 are selected from H or an alkyl group having from about 1 to about 3 carbon atoms; and X is as described previously.

Alternatively, other useful cationic surfactants include amino-amides, wherein in the above structure R_1 is alternatively $R_5CO-(CH_2)_n-$, wherein R_5 is an alkyl group having from about 12 to about 22 carbon atoms, and n is an integer from about 2 to about 6, more preferably from about 2 to about 4, and most preferably from about 2 to about 3. Nonlimiting examples of these cationic emulsifiers include stearamidopropyl PG-dimonium chloride phosphate, stearamidopropyl ethyldimonium ethosulfate, stearamidopropyl dimethyl (myristyl acetate) ammonium chloride, stearamidopropyl dimethyl cetearyl ammonium tosylate, stearamidopropyl dimethyl ammonium chloride, stearamidopropyl dimethyl ammonium lactate, and mixtures thereof.

Nonlimiting examples of quaternary ammonium salt cationic surfactants include those selected from the group consisting of cetyl ammonium chloride, cetyl ammonium bromide, lauryl ammonium chloride, lauryl ammonium bromide, stearyl ammonium chloride, stearyl ammonium bromide, cetyl dimethyl ammonium chloride, cetyl dimethyl ammonium bromide, lauryl dimethyl ammonium chloride, lauryl dimethyl ammonium bromide, stearyl dimethyl ammonium chloride, stearyl dimethyl ammonium bromide, cetyl trimethyl ammonium chloride, cetyl trimethyl ammonium bromide, lauryl trimethyl ammonium chloride, lauryl trimethyl ammonium bromide, stearyl trimethyl ammonium chloride, stearyl trimethyl ammonium bromide, lauryl dimethyl ammonium chloride, stearyl dimethyl cetyl ditallow dimethyl ammonium chloride, dicetyl ammonium chloride, dicetyl ammonium bromide, dilauryl ammonium chloride, dilauryl ammonium bromide, distearyl ammonium chloride, distearyl ammonium bromide, dicetyl methyl ammonium chloride, dicetyl methyl ammonium bromide, dilauryl methyl ammonium chloride, dilauryl methyl ammonium bromide, distearyl methyl ammonium chloride, distearyl dimethyl ammonium chloride, distearyl methyl ammonium bromide, and mixtures thereof. Additional

quaternary ammonium salts include those wherein the C12 to C22 alkyl carbon chain is derived from a tallow fatty acid or from a coconut fatty acid. The term "tallow" refers to an alkyl group derived from tallow fatty acids (usually hydrogenated tallow fatty acids), which generally have mixtures of alkyl chains in the C16 to C18 range. The term "coconut" refers to an alkyl group
5 derived from a coconut fatty acid, which generally have mixtures of alkyl chains in the C12 to C14 range. Examples of quaternary ammonium salts derived from these tallow and coconut sources include ditallow dimethyl ammonium chloride, ditallow dimethyl ammonium methyl sulfate, di(hydrogenated tallow) dimethyl ammonium chloride, di(hydrogenated tallow) dimethyl ammonium acetate, ditallow dipropyl ammonium phosphate, ditallow dimethyl ammonium nitrate,
10 di(coconutalkyl)dimethyl ammonium chloride, di(coconutalkyl)dimethyl ammonium bromide, tallow ammonium chloride, coconut ammonium chloride, stearamidopropyl PG-dimonium chloride phosphate, stearamidopropyl ethyldimonium ethosulfate, stearamidopropyl dimethyl (myristyl acetate) ammonium chloride, stearamidopropyl dimethyl cetearyl ammonium tosylate, stearamidopropyl dimethyl ammonium chloride, stearamidopropyl dimethyl ammonium lactate, and
15 mixtures thereof.

Preferred cationic surfactants useful herein include those selected from the group consisting of dilauryl dimethyl ammonium chloride, distearyl dimethyl ammonium chloride, dimyristyl dimethyl ammonium chloride, dipalmityl dimethyl ammonium chloride, distearyl dimethyl ammonium chloride, and mixtures thereof.

20 The compositions of the present invention can comprise from about 0.1% to about 15%, more preferably from about 0.2% to about 10%, and most preferably from about 0.5% to about 5% of a nonionic surfactant.


Nonlimiting examples of nonionic surfactants for use in the compositions of the present invention are disclosed in McCutcheon's, Detergents and Emulsifiers, North American edition
25 (1986), published by allured Publishing Corporation; and McCutcheon's, Functional Materials, North American Edition (1992); both of which are incorporated by reference herein in their entirety.

Among the nonionic surfactants that are useful herein are those that can be broadly defined as condensation products of long chain alcohols, e.g. C8-30 alcohols, with sugar or starch polymers,
30 i.e., glycosides. These compounds can be represented by the formula $(S)_n-O-R$ wherein S is a sugar moiety such as glucose, fructose, mannose, and galactose; n is an integer of from about 1 to about 1000, and R is a C8-30 alkyl group. Examples of long chain alcohols from which the alkyl group can be derived include decyl alcohol, cetyl alcohol, stearyl alcohol, lauryl alcohol, myristyl alcohol, oleyl alcohol, and the like. Preferred examples of these surfactants include those wherein S is a
35 glucose moiety, R is a C8-20 alkyl group, and n is an integer of from about 1 to about 9. Commercially available examples of these surfactants include decyl polyglucoside (available as

APG 325 CS from Henkel) and lauryl polyglucoside (available as APG 600CS and 625 CS from Henkel).

Other useful nonionic surfactants include the condensation products of alkylene oxides with fatty acids (i.e. alkylene oxide esters of fatty acids). These materials have the general formula $\text{RCO(X)}_n\text{OH}$ wherein R is a C10-30 alkyl group, X is $-\text{OCH}_2\text{CH}_2-$ (i.e. derived from ethylene glycol or oxide) or $-\text{OCH}_2\text{CHCH}_3-$ (i.e. derived from propylene glycol or oxide), and n is an integer from about 1 to about 100. Other nonionic surfactants are the condensation products of alkylene oxides with 2 moles of fatty acids (i.e. alkylene oxide diesters of fatty acids). These materials have the general formula $\text{RCO(X)}_n\text{OOCR}$ wherein R is a C10-30 alkyl group, X is $-\text{OCH}_2\text{CH}_2-$ (i.e. derived from ethylene glycol or oxide) or $-\text{OCH}_2\text{CHCH}_3-$ (i.e. derived from propylene glycol or oxide), and n is an integer from about 1 to about 100. Other nonionic surfactants are the condensation products of alkylene oxides with fatty alcohols (i.e. alkylene oxide ethers of fatty alcohols). These materials have the general formula $\text{R(X)}_n\text{OR'}$ wherein R is a C10-30 alkyl group, X is $-\text{OCH}_2\text{CH}_2-$ (i.e. derived from ethylene glycol or oxide) or $-\text{OCH}_2\text{CHCH}_3-$ (i.e. derived from propylene glycol or oxide), and n is an integer from about 1 to about 100 and R' is H or a C10-30 alkyl group. Still other nonionic surfactants are the condensation products of alkylene oxides with both fatty acids and fatty alcohols [i.e. wherein the polyalkylene oxide portion is esterified on one end with a fatty acid and etherified (i.e. connected via an ether linkage) on the other end with a fatty alcohol]. These materials have the general formula $\text{RCO(X)}_n\text{OR'}$ wherein R and R' are C10-30 alkyl groups, X is $-\text{OCH}_2\text{CH}_2-$ (i.e. derived from ethylene glycol or oxide) or $-\text{OCH}_2\text{CHCH}_3-$ (derived from propylene glycol or oxide), and n is an integer from about 1 to about 100. Nonlimiting examples of these alkylene oxide derived nonionic surfactants include ceteth-1, ceteth-2, ceteth-6, ceteth-10, ceteth-12, cetareth-2, cetareth-6, cetareth-10, cetareth-12, steareth-1, steareth-2, steareth-6, steareth-10, steareth-12, PEG-2 stearate, PEG-4 stearate, PEG-6 stearate, PEG-10 stearate, PEG-12 stearate, steareth-21, PEG-20 glyceryl stearate, PEG-80 glyceryl tallowate, PPG-10 glyceryl stearate, PEG-30 glyceryl cocoate, PEG-80 glyceryl cocoate, PEG-200 glyceryl tallowate, PEG-8 dilaurate, PEG-10 distearate, and mixtures thereof.

Still other useful nonionic surfactants include polyhydroxy fatty acid amide surfactants corresponding to the structural formula:



wherein: R¹ is H, C₁-C₄ alkyl, 2-hydroxyethyl, 2-hydroxypropyl, preferably C₁-C₄ alkyl, more preferably methyl or ethyl, most preferably methyl; R² is C₅-C₃₁ alkyl or alkenyl, preferably C₇-C₁₉ alkyl or alkenyl, more preferably C₉-C₁₇ alkyl or alkenyl, most preferably C₁₁-C₁₅ alkyl

or alkenyl; and Z is a polhydroxyhydrocarbyl moiety having a linear hydrocarbyl chain with a least 3 hydroxyls directly connected to the chain, or an alkoxylated derivative (preferably ethoxylated or propoxylated) thereof. Z preferably is a sugar moiety selected from the group consisting of glucose, fructose, maltose, lactose, galactose, mannose, xylose, and mixtures thereof. An especially preferred surfactant corresponding to the above structure is coconut alkyl N-methyl glucoside amide (i.e., wherein the R^2CO- moiety is derived from coconut oil fatty acids). Processes for making compositions containing polyhydroxy fatty acid amides are disclosed, for example, in G.B. Patent Specification 809,060, published February 18, 1959, by Thomas Hedley & Co., Ltd.; U.S. Patent No. 2,965,576, to E.R. Wilson, issued December 20, 1960; U.S. Patent No. 2,703,798, to A.M. Schwartz, issued March 8, 1955; and U.S. Patent No. 1,985,424, to Piggott, issued December 25, 1934; which are incorporated herein by reference in their entirety.

HUMECTANTS AND MOISTURIZERS

The compositions of the present invention can also comprise one or more humectants or moisturizers. A variety of these materials can be employed and each can be present at a level of from about 0.1% to about 20%, more preferably from about 0.5% to about 10%, and most preferably from about 1% to about 5%. Nonlimiting examples of humectants include materials selected from the group consisting of guanidine; glycolic acid and glycolate salts (e.g., ammonium and quaternary alkyl ammonium); lactic acid and lactate salts (e.g., ammonium and quaternary alkyl ammonium); aloe vera in any of its variety of forms (e.g., aloe vera gel); polyhydroxy alcohols such as sorbitol, glycerol, hexanetriol, propylene glycol, butylene glycol, hexylene glycol and the like; polyethylene glycols; sugars and starches; sugar and starch derivatives (e.g., alkoxylated glucose); hyaluronic acid; lactamide monoethanolamine; acetamide monoethanolamine; and mixtures thereof.

Also, useful are propoxylated glycerols as described in U.S. Patent No. 4,976,953, to Orr et al., issued December 11, 1990, which is incorporated by reference herein in its entirety.

An especially preferred humectant for use herein is glycerol.

INSOLUBLE PARTICLES

The compositions of the present invention can comprise from about 0.1% to about 20%, more preferably from about 0.25% to about 15%, and most preferably from about 0.5% to about 10%, based on the weight of the total composition, of insoluble particles which are useful for enhancing the cleansing effect, when the compositions of the present invention are in the form of a cleansing composition.

The term "insoluble", as used herein, means that the particles are essentially insoluble in the compositions of the present invention. In particular, the insoluble particles should have a solubility less than about 1 gram per 100 grams of composition at 25°C, preferably less than about 0.5 grams per 100 grams of composition at 25°C, and more preferably less than about 0.1 grams per 100 grams of composition at 25°C.

Useful herein are both micronized and conventional size insoluble particles. The micronized particles, for the most part, are of a size that is below the tactile threshold and are essentially nonabrasive to the skin. The conventional size particles are tactilely perceptible and are added for the scrubbing and abrasive effect which they provide.

5 The micronized particles have a mean particle size diameter and particle size distribution such that they are below the tactile perception threshold of most users, and yet are not so small as to be ineffective for aiding in oil, dirt, and debris (e.g., make-up) removal. It is found herein that particles having a mean particle size diameter greater than about 75 microns are tactilely perceived during the cleansing process and it is important to minimize the amount of these larger particles if it is desired that the particles not be felt by the user. Conversely, it is found that particles having a mean particle size diameter of less than about 1 to about 5 microns are generally less effective for providing a cleansing benefit. Without being limited by theory, it is believed that the micronized cleansing particles should be of a size that is on the order of the thickness of the dirt, oil, or debris layer to be cleaned away. This layer is believed to be on the order of a few microns in thickness in most instances. It is therefore found in the present invention that the micronized particles should have a mean particle size diameter from about 1 to about 75 microns, more preferably from about 15 to about 60 microns, and most preferably from about 20 to about 50 microns, so as to provide effective cleansing without being tactilely perceptible. Particles having a wide range of shapes, surface characteristics, and hardness characteristics can be utilized herein provided the particle size requirements are met. Micronized particles of the present invention can be derived from a wide variety of materials including those derived from inorganic, organic, natural, and synthetic sources. Nonlimiting examples of these materials include those selected from the group consisting of almond meal, alumina, aluminum oxide, aluminum silicate, apricot seed powder, attapulgite, barley flour, bismuth oxychloride, boron nitride, calcium carbonate, calcium phosphate, calcium pyrophosphate, calcium sulfate, cellulose, chalk, chitin, clay, corn cob meal, corn cob powder, corn flour, corn meal, corn starch, diatomaceous earth, dicalcium phosphate, dicalcium phosphate dihydrate, fullers earth, hydrated silica, hydroxyapatite, iron oxide, jojoba seed powder, kaolin, loofah, magnesium trisilicate, mica, microcrystalline cellulose, montmorillonite, oat bran, oat flour, oatmeal, peach pit powder, pecan shell powder, polybutylene, polyethylene, polyisobutylene, polymethylstyrene, polypropylene, polystyrene, polyurethane, nylon, teflon (i.e. polytetrafluoroethylene), polyhalogenated olefins, pumice rice bran, rye flour, sericite, silica, silk, sodium bicarbonate, sodium silicoaluminate, soy flour synthetic hectorite, talc, tin oxide, titanium dioxide, tricalcium phosphate, walnut shell powder, wheat bran, wheat flour, wheat starch, zirconium silicate, and mixtures thereof. Also useful are micronized particles made from mixed polymers (e.g., copolymers, terpolymers, etc.), such as polyethylene/polypropylene copolymer, polyethylene/propylene/isobutylene copolymer, polyethylene/styrene copolymer, and the like. Typically, the polymeric and mixed polymeric particles are treated via an oxidation process to destroy impurities

and the like. The polymeric and mixed polymeric particles can also optionally be crosslinked with a variety of common crosslinking agents, nonlimiting examples of which include butadiene, divinyl benzene, methylenebisacrylamide, allyl ethers of sucrose, allyl ethers of pentaerythritol, and mixtures thereof. Other examples of useful micronized particles include waxes and resins such as paraffins, carnuba wax, ozokerite wax, candellila wax, urea-formaldehyde resins, and the like. When such waxes and resins are used herein it is important that these materials are solids at ambient and skin temperatures.

Among the preferred water-insoluble, micronized particulate materials useful herein are the synthetic polymeric particles selected from the group consisting of polybutylene, polyethylene, polyisobutylene, polymethylstyrene, polypropylene, polystyrene, polyurethane, nylon, teflon, and mixtures thereof. Most preferred are polyethylene and polypropylene micronized particles, with the oxidized versions of these materials being especially preferred. Examples of commercially available particles useful herein include the ACumistTM micronized polyethylene waxes available from Allied Signal (Morristown, NJ) available as the A, B, C, and D series in a variety of average particle sizes ranging from 5 microns to 60 microns. Preferred are the ACumistTM A-25, A-30, and A-45 oxidized polyethylene particles having a mean particle size of 25, 30, and 45 microns, respectively. Examples of commercially available polypropylene particles include the Propyltex series available from Micro Powders (Dartek).

The conventional size insoluble particles are well-known to formulation chemists in the art. These particles typically have larger particle sizes than the micronized particles described herein. These particles generally have an average particle size diameter that is about 75 microns or greater, which is about the tactile threshold described above. These conventional size particles typically have average particle sizes ranging up to about 400 microns and larger. These particles can be made from the same materials as for the micronized particles just described. Among the preferred conventional size particulate materials useful herein are the synthetic polymeric particles selected from the group consisting of polybutylene, polyethylene, polyisobutylene, polymethylstyrene, polypropylene, polystyrene, polyurethane, nylon, teflon, and mixtures thereof. Most preferred are polyethylene and polypropylene micronized particles, with the oxidized versions of these materials being especially preferred. An example of a commercially available conventional size particle useful herein is ACuscrubTM 51, available from Allied Signal (Morristown, NJ) having a mean particle size of about 125 microns.

EMULSIFIERS

The compositions herein can comprise various emulsifiers. These emulsifiers are useful for emulsifying the various carrier components of the compositions herein. Suitable emulsifiers can include any of a wide variety of nonionic, cationic, anionic, and zwitterionic emulsifiers disclosed in the prior patents and other references. See McCutcheon's, Detergents and Emulsifiers, North American Edition (1986), published by Allured Publishing Corporation; U.S. Patent No. 5,011,681

to Ciotti et al., issued April 30, 1991; U.S. Patent No. 4,421,769 to Dixon et al., issued December 20, 1983; and U.S. Patent No. 3,755,560 to Dickert et al., issued August 28, 1973; these four references are incorporated herein by reference in their entirety.

Suitable emulsifier types include esters of glycerin, esters of propylene glycol, fatty acid
5 esters of polyethylene glycol, fatty acid esters of polypropylene glycol, esters of sorbitol, esters of sorbitan anhydrides, carboxylic acid copolymers, esters and ethers of glucose, ethoxylated ethers, ethoxylated alcohols, alkyl phosphates, polyoxyethylene fatty ether phosphates, fatty acid amides, acyl lactylates, soaps and mixtures thereof.

Suitable emulsifiers can include, but are not limited to, polyethylene glycol 20 sorbitan
10 monolaurate (polysorbate 20), polyethylene glycol 5 soya sterol, steareth-2, steareth-20, steareth-21, cetareth-20, PPG-2 methyl glucose ether distearate, ceteth-10, polysorbate 80, cetyl phosphate, potassium cetyl phosphate, diethanolamine cetyl phosphate, polysorbate 60, glyceryl stearate, PEG-100 stearate, and mixtures thereof.

The emulsifiers can be used individually or as a mixture of two or more and can comprise
15 from about 0.1% to about 10%, more preferably from about 1% to about 7%, and most preferably from about 1% to about 5% of the compositions of the present invention.

OILS

The compositions of the present invention can comprise from about 0.25% to about 40%,
preferably from about 0.5% to about 25%, and more preferably from about 0.75% to about 15% of
20 an oil selected from the group consisting of mineral oil, petrolatum, C7-C40 branched chain hydrocarbons, C1-C30 alcohol esters of C1-C30 carboxylic acids, C1-C30 alcohol esters of C2-C30 dicarboxylic acids, monoglycerides of C1-C30 carboxylic acids, diglycerides of C1-C30 carboxylic acids, triglycerides of C1-C30 carboxylic acids, ethylene glycol monoesters of C1-C30 carboxylic acids, ethylene glycol diesters of C1-C30 carboxylic acids, propylene glycol monoesters of C1-C30
25 carboxylic acids, propylene glycol diesters of C1-C30 carboxylic acids, C1-C30 carboxylic acid monoesters and polyesters of sugars, polydialkylsiloxanes, polydiarylsiloxanes, polyalkarylsiloxanes, cyclomethicones having 3 to 9 silicon atoms, vegetable oils, hydrogenated vegetable oils, polypropylene glycols, polypropylene glycol C4-C20 alkyl ethers, di C8-C30 alkyl ethers, and mixtures thereof.

30 The oil materials generally having low solubility in water, generally less than about 1% by weight at 25°C. Nonlimiting examples of suitable oil components include, but are not limited to, the following materials. Some of these materials are further described in U.S. Patent No. 4,919,934, to Deckner et al., issued April 24 1990, which is incorporated herein by reference in its entirety.

Mineral oil, which is also known as petrolatum liquid, is a mixture of liquid hydrocarbons
35 obtained from petroleum. See The Merck Index, Tenth Edition, Entry 7048, p. 1033 (1983) and International Cosmetic Ingredient Dictionary, Fifth Edition, vol. 1, p.415-417 (1993), which are incorporated by reference herein in their entirety.

Petrolatum, which is also known as petroleum jelly, is a colloidal system of nonstraight-chain solid hydrocarbons and high-boiling liquid hydrocarbons, in which most of the liquid hydrocarbons are held inside the micelles. See The Merck Index, Tenth Edition, Entry 7047, p. 1033 (1983); Schindler, Drug. Cosmet. Ind., 89, 36-37, 76, 78-80, 82 (1961); and International
5 Cosmetic Ingredient Dictionary, Fifth Edition, vol. 1, p. 537 (1993), which are incorporated by reference herein in their entirety.

Straight and branched chain hydrocarbons having from about 7 to about 40 carbon atoms are useful herein. Nonlimiting examples of these hydrocarbon materials include dodecane, isododecane, squalane, cholesterol, hydrogenated polyisobutylene, docosane (i.e. a C₂₂
10 hydrocarbon), hexadecane, isohexadecane (a commercially available hydrocarbon sold as Permethyl[®] 101A by Presperse, South Plainfield, NJ). Also useful are the C7-C40 isoparaffins, which are C7-C40 branched hydrocarbons.

Useful oils include C1-C30 alcohol esters of C1-C30 carboxylic acids and of C2-C30 dicarboxylic acids, including straight and branched chain materials as well as aromatic derivatives.
15 Also useful are esters such as monoglycerides of C1-C30 carboxylic acids, diglycerides of C1-C30 carboxylic acids, triglycerides of C1-C30 carboxylic acids, ethylene glycol monoesters of C1-C30 carboxylic acids, ethylene glycol diesters of C1-C30 carboxylic acids, propylene glycol monoesters of C1-C30 carboxylic acids, and propylene glycol diesters of C1-C30 carboxylic acids. Straight chain, branched chain and aryl carboxylic acids are included herein. Also useful are propoxylated
20 and ethoxylated derivatives of these materials. Nonlimiting examples include diisopropyl sebacate, diisopropyl adipate, isopropyl myristate, isopropyl palmitate, myristyl propionate, ethylene glycol distearate, 2-ethylhexyl palmitate, isodecyl neopentanoate, C₁₂₋₁₅ alcohols benzoate, di-2-ethylhexyl maleate, cetyl palmitate, myristyl myristate, stearyl stearate, cetyl stearate, behenyl behenrate, dioctyl maleate, dioctyl sebacate, diisopropyl adipate, cetyl octanoate, diisopropyl
25 dilinoleate, caprylic/capric triglyceride, PEG-6 caprylic/capric triglyceride, PEG-8 caprylic/capric triglyceride, and mixtures thereof.

Also useful are various C1-C30 monoesters and polyesters of sugars and related materials. These esters are derived from a sugar or polyol moiety and one or more carboxylic acid moieties. Depending on the constituent acid and sugar, these esters can be in either liquid or solid form at
30 room temperature. Examples of liquid esters include: glucose tetraoleate, the glucose tetraesters of soybean oil fatty acids (unsaturated), the mannose tetraesters of mixed soybean oil fatty acids, the galactose tetraesters of oleic acid, the arabinose tetraesters of linoleic acid, xylose tetralinoleate, galactose pentaoleate, sorbitol tetraoleate, the sorbitol hexaesters of unsaturated soybean oil fatty acids, xylitol pentaoleate, sucrose tetraoleate, sucrose pentaoleate, sucrose hexaoleate, sucrose
35 hepatoleate, sucrose octaoleate, and mixtures thereof. Examples of solid esters include: sorbitol hexaester in which the carboxylic acid ester moieties are palmitoleate and arachidate in a 1:2 molar ratio; the octaester of raffinose in which the carboxylic acid ester moieties are linoleate and behenate

in a 1:3 molar ratio; the heptaester of maltose wherein the esterifying carboxylic acid moieties are sunflower seed oil fatty acids and lignocerate in a 3:4 molar ratio; the octaester of sucrose wherein the esterifying carboxylic acid moieties are oleate and behenate in a 2:6 molar ratio; and the octaester of sucrose wherein the esterifying carboxylic acid moieties are laurate, linoleate and behenate in a 1:3:4 molar ratio. A preferred solid material is sucrose polyester in which the degree of esterification is 7-8, and in which the fatty acid moieties are C18 mono- and/or di-unsaturated and behenic, in a molar ratio of unsaturates:behenic of 1:7 to 3:5. A particularly preferred solid sugar polyester is the octaester of sucrose in which there are about 7 behenic fatty acid moieties and about 1 oleic acid moiety in the molecule. The ester materials are further described in, U.S. Patent No. 2,831,854, U.S. Patent No. 4,005,196, to Jandacek, issued January 25, 1977; U.S. Patent No. 4,005,195, to Jandacek, issued January 25, 1977, U.S. Patent No. 5,306,516, to Letton et al., issued April 26, 1994; U.S. Patent No. 5,306,515, to Letton et al., issued April 26, 1994; U.S. Patent No. 5,305,514, to Letton et al., issued April 26, 1994; U.S. Patent No. 4,797,300, to Jandacek et al., issued January 10, 1989; U.S. Patent No. 3,963,699, to Rizzi et al, issued June 15, 1976; U.S. Patent No. 4,518,772, to Volpenhein, issued May 21, 1985; and U.S. Patent No. 4,517,360, to Volpenhein, issued May 21, 1985; all of which are incorporated by reference herein in their entirety.

Silicones such as polydialkylsiloxanes, polydiarylsiloxanes, polyalkarylsiloxanes, and cyclomethicones having 3 to 9 silicon atoms are useful oils. These silicones include both volatile and nonvolatile materials. These silicones are disclosed in U.S. Patent No. 5,069,897, to Orr, issued December 3, 1991, which is incorporated by reference herein in its entirety. The polyalkylsiloxanes include, for example, polyalkylsiloxanes with viscosities of from about 0.5 to about 100,000 centistokes at 25°C. Such polyalkylsiloxanes correspond to the general chemical formula $R_3SiO[R_2SiO]_xSiR_3$ wherein R is an alkyl group (preferably R is methyl or ethyl, more preferably methyl) and x is an integer from 0 to about 500, chosen to achieve the desired molecular weight. Commercially available polyalkylsiloxanes include the polydimethylsiloxanes, which are also known as dimethicones, nonlimiting examples of which include the Vicasil[®] series sold by General Electric Company and the Dow Corning[®] 200 series sold by Dow Corning Corporation. Specific examples of polydimethylsiloxanes useful as emollients herein include Dow Corning[®] 200 fluid having a viscosity of 0.65 centistokes and a boiling point of 100°C, Dow Corning[®] 225 fluid having a viscosity of 10 centistokes and a boiling point greater than 200°C, and Dow Corning[®] 200 fluids having viscosities of 50, 350, and 12,500 centistokes, respectively, and boiling points greater than 200°C. Cyclic polyalkylsiloxanes useful herein include those corresponding to the general chemical formula $[SiR_2-O]_n$ wherein R is an alkyl group (preferably R is methyl or ethyl, more preferably methyl) and n is an integer from about 3 to about 9, more preferably n is an integer from about 3 to about 7, and most preferably n is an integer from about 4 to about 6. When R is methyl, these materials are typically referred to as cyclomethicones. Commercially available cyclomethicones include Dow Corning[®] 244 fluid having a viscosity of 2.5 centistokes, and a boiling point of 172°C,

which primarily contains the cyclomethicone tetramer (i.e. n=4), Dow Corning® 344 fluid having a viscosity of 2.5 centistokes and a boiling point of 178°C, which primarily contains the cyclomethicone pentamer (i.e. n=5), Dow Corning® 245 fluid having a viscosity of 4.2 centistokes and a boiling point of 205°C, which primarily contains a mixture of the cyclomethicone tetramer and pentamer (i.e. n=4 and 5), and Dow Corning® 345 fluid having a viscosity of 4.5 centistokes and a boiling point of 217°, which primarily contains a mixture of the cyclomethicone tetramer, pentamer, and hexamer (i.e. n=4, 5, and 6). Also useful are materials such as trimethylsiloxysilicate, which is a polymeric material corresponding to the general chemical formula $[(CH_2)_3SiO_{1/2}]_x[SiO_2]_y$, wherein x is an integer from about 1 to about 500 and y is an integer from about 1 to about 500. A commercially available trimethylsiloxysilicate is sold as a mixture with dimethicone as Dow Corning® 593 fluid. Also useful herein are dimethiconols, which are hydroxy terminated dimethyl silicones. These materials can be represented by the general chemical formulas $R_3SiO[R_2SiO]_xSiR_2OH$ and $HOR_2SiO[R_2SiO]_xSiR_2OH$ wherein R is an alkyl group (preferably R is methyl or ethyl, more preferably methyl) and x is an integer from 0 to about 500, chosen to achieve the desired molecular weight. Commercially available dimethiconols are typically sold as mixtures with dimethicone or cyclomethicone (e.g. Dow Corning® 1401, 1402, and 1403 fluids). Also useful herein are polyalkylaryl siloxanes, with polymethylphenyl siloxanes having viscosities from about 15 to about 65 centistokes at 25°C being preferred. These materials are available, for example, as SF 1075 methylphenyl fluid (sold by General Electric Company) and 556 Cosmetic Grade phenyl trimethicone fluid (sold by Dow Corning Corporation).

Vegetable oils and hydrogenated vegetable oils are also useful herein. Examples of vegetable oils and hydrogenated vegetable oils include safflower oil, castor oil, coconut oil, cottonseed oil, menhaden oil, palm kernel oil, palm oil, peanut oil, soybean oil, rapeseed oil, linseed oil, rice bran oil, pine oil, sesame oil, sunflower seed oil, hydrogenated safflower oil, hydrogenated castor oil, hydrogenated coconut oil, hydrogenated cottonseed oil, hydrogenated menhaden oil, hydrogenated palm kernel oil, hydrogenated palm oil, hydrogenated peanut oil, hydrogenated soybean oil, hydrogenated rapeseed oil, hydrogenated linseed oil, hydrogenated rice bran oil, hydrogenated sesame oil, hydrogenated sunflower seed oil, and mixtures thereof.

Also useful are polypropylene glycols, C4-C20 alkyl ethers of polypropylene glycols, C1-C20 carboxylic acid esters of polypropylene glycols, and di-C8-C30 alkyl ethers. Nonlimiting examples of these materials include PPG-14 butyl ether, PPG-15 stearyl ether, PPG-9, PPG-12, PPG-15, PPG-17, PPG-20, PPG-26, PPG-30, PPG-34, dioctyl ether, dodecyl octyl ether, and mixtures thereof.

SUNSCREENS AND TANNING AGENTS

Also useful herein are sunscreens. A wide variety of sunscreens are described in U.S. Patent No. 5,087,445, to Haffey et al., issued February 11, 1992; U.S. Patent No. 5,073,372, to Turner et al., issued December 17, 1991; U.S. Patent No. 5,073,371, to Turner et al. issued December 17, 1991; and Segarin, et al., at Chapter VIII, pages 189 et seq., of Cosmetics

Science and Technology, all of which are incorporated herein by reference in their entirety. Nonlimiting examples of sunscreens which are useful in the compositions of the present invention are those selected from the group consisting of 2-ethylhexyl *p*-methoxycinnamate, 2-ethylhexyl *N,N*-dimethyl-*p*-aminobenzoate, *p*-aminobenzoic acid, 2-phenylbenzimidazole-5-sulfonic acid, 5 octocrylene, oxybenzone, homomenthyl salicylate, octyl salicylate, 4,4'-methoxy-*t*-butyldibenzoylmethane, 4-isopropyl dibenzoylmethane, 3-benzylidene camphor, 3-(4-methylbenzylidene) camphor, titanium dioxide, zinc oxide, silica, iron oxide, and mixtures thereof. Still other useful sunscreens are those disclosed in U.S. Patent No. 4,937,370, to Sabatelli, issued June 26, 1990; and U.S. Patent No. 4,999,186, to Sabatelli et al., issued March 12, 1991; these two 10 references are incorporated by reference herein in their entirety. The suncreening agents disclosed therein have, in a single molecule, two distinct chromophore moieties which exhibit different ultra-violet radiation absorption spectra. One of the chromophore moieties absorbs predominantly in the UVB radiation range and the other absorbs strongly in the UVA radiation range. These sunscreening agents provide higher efficacy, broader UV absorption, lower skin penetration and 15 longer lasting efficacy relative to conventional sunscreens. Especially preferred examples of these sunscreens include those selected from the group consisting of 4-*N,N*-(2-ethylhexyl)methylaminobenzoic acid ester of 2,4-dihydroxybenzophenone, 4-*N,N*-(2-ethylhexyl)methylaminobenzoic acid ester with 4-hydroxydibenzoylmethane, 4-*N,N*-(2-ethylhexyl)methylaminobenzoic acid ester of 2-hydroxy-4-(2-hydroxyethoxy)benzophenone, 4-*N,N*-(2-ethylhexyl)-methylaminobenzoic acid ester 20 of 4-(2-hydroxyethoxy)dibenzoylmethane, and mixtures thereof. Generally, the sunscreens can comprise from about 0.5% to about 20% of the compositions useful herein. Exact amounts will vary depending upon the sunscreen chosen and the desired Sun Protection Factor (SPF). SPF is a commonly used measure of photoprotection of a sunscreen against erythema. See Federal Register, Vol. 43, No. 166, pp. 38206-38269, August 25, 1978, which is incorporated herein by reference in 25 its entirety.

Examples of artificial tanning agents and accelerators include dihydroxyacetone, tyrosine, tyrosine esters such as ethyl tyrosinate, and phospho-DOPA.

METHOD OF FORMING THE COMPLEX

The complex that is believed to be formed between the amphoteric and anionic surfactants 30 of the present invention is preferably preformed and added to the active ingredient and any remaining components of the compositions of the present invention. The amphoteric and anionic surfactants are preferably combined in aqueous solution, thereby forming what is believed to be an aqueous dispersion of the complex.

METHODS OF TREATING THE SKIN

35 The present invention also relates to methods wherein an effective amount of the composition of the present invention is applied to the skin. These compositions are useful for delivering active ingredients to the skin. A wide range of quantities of the compositions of the

present invention can be used. Quantities which are typically applied can range from about 0.1 mg/cm² to about 25 mg/cm².

In preferred embodiments, the compositions of the present invention are also useful for personal cleansing, especially for cleansing of the face and neck areas. Typically, a suitable or effective amount of the cleansing composition is applied to the area to be cleansed. Alternatively, a suitable amount of the cleansing composition can be applied via intermediate application to a washcloth, sponge, pad, cotton ball or other application device. If desired, the area to be cleansed can be premoistened with water. It has been found that the compositions of the present invention can be combined with water during the cleansing process and rinsed-off from the skin. Alternatively, the composition can be used alone and wiped-off from the skin using a pad, cotton ball, tissue, or other like device. The cleansing process is typically a two-step process involving application of the composition followed either by rinsing of the product with water or wiping without the use of water. Generally, an effective amount of composition to be used will depend upon the needs and usage habits of the individual.

EXAMPLES

The following examples further describe and demonstrate embodiments within the scope of the present invention. The examples are given solely for the purpose of illustration and are not to be construed as limitations of the present invention, as many variations thereof are possible without departing from the spirit and scope of the invention.

Ingredients are identified by chemical or CTFA name.

Example 1

A leave-on lotion composition containing salicylic acid is prepared by combining the following ingredients using conventional mixing techniques.

25

	<u>Ingredients</u>	<u>Weight Percent</u>
	<u>Phase A</u>	
	Water	QS 100
30	Glycerin	3.00
	Tetrasodium EDTA	0.02
	<u>Phase B</u>	
	PPG-15 Stearyl Ether	4.00
35	Stearyl Alcohol	0.75
	Salicylic Acid	2.00
	Cetyl Alcohol	0.75

	Stearth-21	0.45
	Stearth-2	0.05
	Dimethicone	0.60
	Polyquaternium-37 (and) Mineral Oil (and) PPG-1 Trideceth-6	1.50

5

Phase C

	Triethanolamine	0.15
--	-----------------	------

Phase D

10	Fragrance	0.10
----	-----------	------

Phase E

	Cetyl Dimethyl Betaine	2.00
	Sodium Lauryl Sulfate	1.00

15

In a suitable vessel, the Phase A ingredients are heated with stirring to about 75°C. In a separate vessel, the Phase B ingredients are heated with stirring to about 75°C. Phase B is then added to Phase A with mixing. Next Phase C is added with mixing. Next the fragrance is added with mixing. Next, the mixture is cooled to 35°C. In a separate vessel, the Phase E ingredients are combined and added to the remaining mixture with stirring.

20

The resulting leave-on composition is useful for preventing and treating acne while being mild to the skin.

Alternatively, a composition is prepared in which the sodium lauryl sulfate is replaced with sodium lauroyl isethionate.

25

Example 2

A personal cleanser composition containing salicylic acid is prepared by combining the following ingredients using conventional mixing techniques.

30

IngredientsWeight PercentPhase A

	Water	QS 100
	Glycerin	3.00
35	Disodium EDTA	0.01

Phase B

	PPG-15 Stearyl Ether	4.00
	Stearyl Alcohol	2.88
	Salicylic Acid	2.00
	Cetyl Alcohol	0.80
5	Distearyl Dimethyl Ammonium Chloride	1.50
	Steareth-21	0.50
	Behenyl Alcohol	0.32
	PPG-30	0.25
	Steareth-2	0.25
10		
	<u>Phase C</u>	
	Oxidized Polyethylene Beads ¹	1.00
	Fragrance	0.27
15	<u>Phase D</u>	
	Cocamidopropyl Betaine	2.00
	Sodium Lauryl Sulfate	1.00

20 ¹Available as AcucscrubTM 51 from Allied Signal Corporation.

In a suitable vessel, the Phase A ingredients are heated with stirring to about 75°C. In a separate vessel, the Phase B ingredients are heated with stirring to about 75°C. Phase B is then added to Phase A with mixing. Next, the oxidized polyethylene beads are added slowly with mixing to prevent agglomeration. Next the fragrance is added with mixing. Next, the mixture is cooled to 35°C. In a separate vessel, the Phase D ingredients are combined and added to the remaining mixture with stirring.

The resulting cleansing composition is useful for cleansing the skin.

Alternatively, a composition is prepared in which the sodium lauryl sulfate is replaced with sodium lauroyl isethionate.

Alternatively, a composition is prepared in which the salicylic acid is replaced with 2.5% benzoyl peroxide, with the water being adjusted accordingly.

Example 3

35 A personal cleanser composition containing salicylic acid and menthol is prepared by combining the following ingredients using conventional mixing techniques.

	<u>Ingredients</u>	<u>Weight Percent</u>
	<u>Phase A</u>	
	Water	QS 100
5	Glycerin	3.00
	Disodium EDTA	0.01
	<u>Phase B</u>	
	PPG-15 Stearyl Ether	4.00
10	Stearyl Alcohol	2.88
	Salicylic Acid	2.00
	Distearyl Dimethyl Ammonium Chloride	1.50
	Cetyl Alcohol	0.80
	Steareth-21	0.50
15	Behenyl Alcohol	0.32
	PPG-30	0.25
	Steareth-2	0.25
	<u>Phase C</u>	
20	Oxidized Polyethylene Beads ¹	1.00
	Fragrance	0.27
	Menthol	0.05
	<u>Phase D</u>	
25	Cetyl Dimethyl Betaine	2.00
	Sodium Lauryl Sulfate	1.00

¹ Available as AcucscrubTM 51 from Allied Signal Corporation.

30

In a suitable vessel, the Phase A ingredients are heated with stirring to about 75°C. In a separate vessel, the Phase B ingredients are heated with stirring to about 75°C. Phase B is then added to Phase A with mixing. Next, the oxidized polyethylene beads are added slowly with mixing to prevent agglomeration. Next the fragrance and menthol are added with mixing. Next, the mixture is cooled to 35°C. In a separate vessel, the Phase D ingredients are combined and added to the remaining mixture with stirring.

35

The resulting cleansing composition is useful for cleansing the skin.

Alternatively, a composition is prepared in which the sodium lauryl sulfate is replaced with sodium lauroyl isetheonate.

Alternatively, a composition is prepared in which the salicylic acid is replaced with 2.5% benzoyl peroxide with the water being adjusted correspondingly.

5

Example 4

A leave-on lotion composition containing benzoyl peroxide is prepared by combining the following ingredients using conventional mixing techniques.

10	<u>Ingredients</u>	<u>Weight Percent</u>
	<u>Phase A</u>	
	Water	QS 100
	Glycerin	4.00
15	Disodium EDTA	0.10
	Carbomer	0.60
	Acrylates/C10-30 Alkylacrylates Crosspolymer	0.05
	<u>Phase B</u>	
20	Stearyl Alcohol	2.25
	Cetyl Alcohol	2.25
	Steareth-100	0.50
	Glyceryl Hydroxystearate	0.74
	Dimethicone	0.60
25	<u>Phase C</u>	
	Triethanolamine	0.50
	<u>Phase D</u>	
30	Benzoyl Peroxide	2.50
	<u>Phase E</u>	
	Cetyl Dimethyl Betaine	1.00
	Sodium Lauryl Sulfate	0.50

35

In a suitable vessel, the Phase A ingredients are heated with stirring to about 75°C. In a separate vessel, the Phase B ingredients are heated with stirring to about 75°C. Phase B is then added to

Phase A with mixing. Next Phase C is added with mixing. Next, the mixture is cooled to 35°C. Next the benzoyl peroxide is added with mixing. In a separate vessel, the Phase E ingredients are combined and added to the remaining mixture with stirring.

5 The resulting leave-on composition is useful for preventing and treating acne while being mild to the skin.

Alternatively, a composition is prepared in which the cetyl dimethyl betaine is replaced with stearyl dimethyl betaine.

Example 5

10 A personal cleansing gel composition containing glycolic acid is prepared by combining the following ingredients using conventional mixing techniques.

	<u>Ingredients</u>	<u>Weight Percent</u>
15	<u>Phase A</u>	
	Water	QS 100
	Glycerin	4.00
	Disodium EDTA	0.10
	Dimethicone	0.20
20	PVM/MA Decadiene Crosspolymer	1.00
	Glycolic Acid	2.00
	Sodium Hydroxide	0.80
	<u>Phase B</u>	
25	Cetyl Dimethyl Betaine	1.00
	Sodium Lauryl Sulfate	0.50

In a suitable vessel, the Phase A ingredients are mixed vigorously. In a separate vessel, the Phase B ingredients are combined and added to the remaining mixture with stirring.

30 The resulting cleansing gel composition is useful for reducing the appearance of pore size while being mild to the skin and cleansing effectively.


Alternatively, a composition is prepared in which glycolic acid is replaced with salicylic acid and the resulting composition is useful for preventing and treating acne and blemishes while being mild to the skin and cleansing effectively.

35 Alternatively, a composition is prepared in which the cetyl dimethyl betaine is replaced with cocamidopropyl betaine.

WHAT IS CLAIMED IS:

1. A topical personal care composition comprising:

(a) from 0.1% to 20% by weight of an amphoteric surfactant having the following structure



wherein R^1 is unsubstituted, saturated or unsaturated, straight or branched chain alkyl having from 9 to 22 carbon atoms; m is an integer from 1 to 3; n is 0 or 1; R^2 and R^3 are independently selected from alkyl having from 1 to 3 carbon atoms and monohydroxyalkyl having from 1 to 3 carbon atoms; R^4 is selected from saturated or unsaturated alkyl having from 1 to 5 carbon atoms and saturated or unsaturated monohydroxyalkyl having from 1 to 5 carbon atoms; X is selected from the group consisting of CO_2 , SO_3 , and SO_4 ; and pharmaceutically acceptable salts of the foregoing compounds;


- 15 (b) from 0.1% to 20% by weight of an anionic surfactant,
 (c) from 0.001% to 20% of an active ingredient, and
 (d) from 40% to 99.799% by weight water.

2. A composition according to Claim 1 wherein for said amphoteric surfactant in (a), R^2 and R^3 are selected from the group consisting of CH_3 , $\text{CH}_2\text{CH}_2\text{OH}$ and $\text{CH}_2\text{CH}_2\text{CH}_2\text{OH}$; X is selected from CO_2 and SO_3 ; and m is 2 or 3; preferably wherein R^2 and R^3 are selected from the group consisting of CH_3 , $\text{CH}_2\text{CH}_2\text{OH}$, and $\text{CH}_2\text{CH}_2\text{CH}_2\text{OH}$; X is selected from the group consisting of CO_2 and SO_3 , and m is 2 or 3; more preferably wherein R^4 has from 1 to 3 carbon atoms when X is CO_2 , and R^4 has from 2 to 4 carbon atoms when X is SO_3 ; and even more preferably wherein R^1 has from 11 to 18 carbon atoms; R^2 and R^3 are CH_3 ; and R^4 has 1 carbon atom when X is CO_2 , and R^4 has 3 carbon atoms when X is SO_3 .

3. A composition according to Claim 2 wherein R^4 has 1 carbon atom, X is CO_2 , m is 3, and n is 1.

4. A composition according to Claim 2 wherein R^4 has 3 carbon atoms, X is SO_3 , m is 3 and n is 1.

5. A composition according to Claim 1 where said amphoteric surfactant is selected from the group consisting of cetyl dimethyl betaine, cocoamidopropyl betaine, stearyl dimethyl betaine, cocoamidopropyl hydroxy sultaine, and mixtures thereof; preferably wherein the amphoteric surfactant is cetyl dimethyl betaine; and wherein said anionic surfactant is selected from the group consisting of sodium lauryl sulfate, sodium cetyl sulfate, ammonium cocoyl isethionate, sodium lauroyl isethionate, sodium lauroyl sarcosinate, and mixtures thereof; preferably wherein said anionic surfactant is sodium lauryl sulfate.
6. A composition according to Claim 1 wherein said active ingredient is selected from the group consisting of selected from the group consisting of salicylic acid, benzoyl peroxide, cis-retinoic acid, retinol, phytic acid, trans-retinoic acid, N-acetyl L-cysteine, azelaic acid, lipoic acid, resorcinol, lactic acid, glycolic acid, ibuprofen, naproxen, hydrocortisone, phenoxyethanol, phenoxypropanol, phenoxyisopropanol, 2,4,4'-trichloro-2'-hydroxy diphenyl ether, 3,4,4'-trichlorocarbanilide, and mixtures thereof; preferably wherein said active ingredient is selected from the group consisting of salicylic acid, benzoyl peroxide, and mixtures thereof.
7. A composition according to Claim 1 wherein said composition further comprises from 0.1% to 20% of a cationic surfactant; preferably wherein said cationic surfactant has the formula:



wherein in this formula for said cationic surfactant, R_1 is an alkyl group having from 12 to 22 carbon atoms, R_2 is H or an alkyl group having from 1 to 22 carbon atoms, R_3 and R_4 are independently selected from H or an alkyl group having from 1 to 3 carbon atoms, and X is an anion selected from chloride, bromide, iodide, acetate, phosphate, nitrate, sulfate, methyl sulfate, ethyl sulfate, tosylate, lactate, citrate, glycolate, and mixtures thereof; more preferably wherein said cationic surfactant is selected from the group consisting of dilauryl dimethyl ammonium chloride, distearyl dimethyl ammonium chloride, dimyristyl dimethyl ammonium chloride, dipalmityl dimethyl ammonium chloride, and mixtures thereof.

8. A composition according to Claim 14 wherein said composition further comprises from 0.1% to 20% of a humectant; preferably wherein said humectant is glycerol.

9. A method for treating skin comprising the step of applying to the skin from 0.5 mg/cm² to 25 mg/cm² of the composition of any of Claims 1-8.
10. A method of preparing a composition according to any of claims 1-8 comprising the steps of:
- (i) combining an aqueous solution of said amphoteric surfactant and an aqueous solution of said anionic surfactant to form an aqueous solution of a complex of said amphoteric and said anionic surfactant, and
 - (ii) combining said complex with said active ingredient.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 96/11789

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 A61K7/50

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	WO,A,96 06595 (THE PROCTER & GAMBLE COMPANY) 7 March 1996 see examples ---	1-8
P,X	WO,A,96 02225 (UNILEVER PLC) 1 February 1996 see the whole document ---	1-7
P,X	WO,A,96 05798 (THE PROCTER & GAMBLE COMPANY) 29 February 1996 see the whole document ---	1-7
X	WO,A,91 17237 (THE PROCTER & GAMBLE COMPANY) 14 November 1991 see the whole document --- -/-	1,6

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- *Z* document member of the same patent family

Date of the actual completion of the international search

27 November 1996

Date of mailing of the international search report

11. 12. 96

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer

Couckuyt, P

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 96/11789

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO,A,95 15150 (THE PROCTER & GAMBLE COMPANY) 8 June 1995 see the whole document ---	1-6
P,X	WO,A,95 24179 (THE PROCTER & GAMBLE COMPANY) 14 September 1995 see the whole document ---	1,5,7
X	CHEMICAL ABSTRACTS, vol. 106, no. 12, 23 March 1987 Columbus, Ohio, US; abstract no. 89959, XP002019636 & JP,A,61 263 910 (SHISEIDO CO. LTD) 21 November 1986 see abstract -----	1-3,5,6

INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No

PCT/US 96/11789

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9606595	07-03-96	AU-A- 3330095	22-03-96
WO-A-9602225	01-02-96	AU-A- 2934195	16-02-96
WO-A-9605798	29-02-96	NONE	
WO-A-9117237	14-11-91	AU-A- 7560091	27-11-91
		CA-A- 2078441	28-10-91
		CN-A- 1055109	09-10-91
		EP-A- 0526473	10-02-93
		NZ-A- 237584	27-09-94
WO-A-9515150	08-06-95	CA-A- 2174633	08-06-95
		EP-A- 0730442	11-09-96
WO-A-9524179	14-09-95	AU-A- 1982595	25-09-95
		CA-A- 2185231	14-09-95